

Dynamics of pairwise motions

Roman Juszkiewicz^{1,3}, Volker Springel², and Ruth Durrer¹

ABSTRACT

We derive a simple closed-form expression, relating $v_{12}(r)$ – the mean relative velocity of pairs of galaxies at fixed separation r – to the two-point correlation function of mass density fluctuations, $\xi(r)$. We compare our analytic model for $v_{12}(r)$ with N-body simulations, and find excellent agreement in the entire dynamical range probed by the simulations ($0.1 \lesssim \xi \lesssim 1000$). Our results can be used to estimate the cosmological density parameter, Ω , directly from redshift-distance surveys, like Mark III.

Subject headings: Cosmology: theory – observation – peculiar velocities: large scale flows

1. An analytical model for $v_{12}(r)$

Most dynamical estimates of the cosmological density parameter, Ω , use the gravitational effect of departures from a strictly homogeneous distribution of mass. One such dynamical estimator can be constructed by using an equation expressing the conservation of particle pairs in a self-gravitating gas. This equation was derived by Davis & Peebles (1977) from the BBGKY theory (see also Peebles 1980, hereafter LSS). Consider a pair of particles at a comoving separation vector $\vec{x}$ and cosmological time t , moving with a mean (pair-weighted) relative velocity $v_{12}(x, t) \vec{x} / x$. Its magnitude is related to the two-point correlation function of density fluctuations, $\xi(x, t)$, by the pair conservation equation (LSS),

$$\frac{a}{3[1 + \xi(x, a)]} \frac{\partial \bar{\xi}(x, a)}{\partial a} = - \frac{v_{12}(x, a)}{Hr}, \quad (1)$$

where $a(t)$ is the expansion factor, $r = ax$ is the proper separation, $H(a)$ is the Hubble parameter, while $\bar{\xi}(x, a)$ is the two-point correlation function, averaged over a ball of comoving radius x :

$$\bar{\xi}(x, a) = 3x^{-3} \int_0^x \xi(y, a) y^2 dy. \quad (2)$$

At the present cosmological time $a = 1$, $x = r$ and $H = 100 \, h \text{ km s}^{-1} \text{ Mpc}^{-1}$, where h is the conventional dimensionless parameterization for the Hubble constant. There are two well known

¹Département de Physique Théorique, Université de Genève, CH-1211 Genève, Switzerland

²Max-Planck-Institut für Astrophysik, D-85740 Garching, Germany

³On leave from Copernicus Astronomical Center, 00-716 Warsaw, Poland

approximate solutions of (1). They are: the small separation limit, where $\xi(x) \gg 1$ (stable clustering regime), and the large separation limit, where $|\xi| \ll 1$ (linear regime). The stable clustering solution is (LSS, §71)

$$v_{12}(x, a) = -Hr, \quad (3)$$

as expected for virialized clusters on sufficiently small scales. The linear solution is given by the first terms in perturbative expansions for v_{12} and ξ , which for the correlation is given by

$$\xi = \xi^{(1)} + \xi^{(2)} + \dots, \quad (4)$$

with $\xi^{(2)} = O[\xi^{(1)}]^2$, etc. The growing mode of the linear solution is

$$\xi^{(1)}(x, a) = \xi_1(x) D(a)^2, \quad (5)$$

where $D(a)$ is the usual linear growth factor (LSS, §11; we neglect the decaying mode). The general technique for deriving $\xi^{(2)}$ and higher order terms for initially Gaussian density fluctuations in an Einstein-de Sitter universe was introduced by Juszkiewicz et al. (1980, 1984), Vishniac (1983) and Fry (1984); their results were recently generalized to a wider class of cosmological models including those considered below (Juszkiewicz et al. 1993, Bouchet et al. 1995). These calculations show that the second order term in the expansion for ξ can be written as

$$\xi^{(2)}(x, a) = \xi_2(x) D(a)^4, \quad (6)$$

where ξ_2 is a function of x alone. Substituting $\xi = D^2 \xi_1 + D^4 \xi_2$ into eq. (1) and solving for v_{12} we get

$$v_{12} = -\frac{2}{3} H r f [\bar{\xi}^{(1)} - \bar{\xi}^{(1)} \xi^{(1)} + 2 \bar{\xi}^{(2)}] + O[\xi^{(1)}]^3, \quad (7)$$

where $f \equiv d \ln D / d \ln a \approx \Omega^{0.6}$ (which is a good approximation if $\Lambda = 0$ or $\Omega + \Omega_\Lambda = 1$; $\Omega_\Lambda \equiv \Lambda / 3H^2$). The bars above ξ -s denote averaging over a ball of comoving radius x . If the logarithmic slope of the correlation function, $\gamma(x) \equiv -d \ln \xi^{(1)}(x, a) / d \ln x$, is a constant in the range $0 \leq \gamma < 2$. Then $\bar{\xi}^{(2)}$ is related to $\bar{\xi}^{(1)}$ by a simple closed-form expression,

$$\bar{\xi}^{(2)}(x, a) = \alpha(\gamma) [\bar{\xi}^{(1)}(x, a)]^2, \quad (8)$$

where α is a constant (Łokas et al. 1996), and the function $\alpha(\gamma)$ can be expressed in terms of Euler's Γ functions (Scoccimarro & Frieman 1996). In the range $0 \leq \gamma < 2$, this expression is well approximated by a fitting formula

$$\alpha = 1.84 - 1.1 \gamma - 0.84 (\gamma/2)^{10}. \quad (9)$$

For $\gamma \geq 2$, perturbation theory diverges, but N-body simulations suggest that the relation (8) remains valid, provided the $\alpha(\gamma)$ dependence for $\gamma \gtrsim 1.5$ is derived from the so-called universal scaling relation (USH) – an empirical formula for the nonlinear power spectrum, extracted from $\gamma \geq 1$ scale-free and cold dark matter (CDM) simulations (Jain et al. 1995, Scoccimarro & Frieman 1996). Since the USH approach fails when $\gamma < 1$, while the perturbation theory (PT) fails for $\gamma \geq 2$, we propose to trade accuracy for an extended range of validity and interpolate the expression

$\alpha(\gamma)$ between the PT formula for $\gamma < 1$ and the USH result for $\gamma > 1$ (Scoccimarro & Frieman 1996, Fig.20), and replace eq. (9) with

$$\alpha \approx 1.2 - 0.65\gamma. \quad (10)$$

The allowed range of slopes for the above expression is $0 < \gamma < 3$. To extend the range of validity of our perturbative solution (7) into the nonlinear regime, we propose the following Ansatz:

$$v_{12}(x, a) = -\frac{2}{3} H r f \bar{\xi}(x, a) \left[1 + \alpha \bar{\xi}(x, a) \right], \quad (11)$$

$$\bar{\bar{\xi}}(x, a) = \bar{\xi}(x, a) [1 + \xi(x, a)]^{-1}. \quad (12)$$

The expression (11) agrees with the perturbative expansion (7) when $\xi \rightarrow 0$; it also approximates the stable clustering limit when $x \rightarrow 0$. Indeed, when ξ is large, $-v_{12}/Hr \approx (2/3)f(\Omega)(1 + \alpha)$ which is of order unity for all reasonable values of Ω and α .

Eq. (11) can be used to predict the relative velocity v_{12} at any separation r . The only input necessary for that is the correlation function in an interval of separations $\leq r$. This is very different from the semi-analytic expression for $v_{12}(r)$, derived by Mo et al. (1997). In their case, the knowledge of the present $\xi(r)$ is not sufficient to calculate $v_{12}(r)$; they use the USH approach and a large array of parameters instead.

2. N-body simulations

In this section we compare our analytic expression for v_{12} with the results from high-resolution AP³M simulations of 256^3 dark matter particles, kindly provided to us by the Virgo collaboration (Jenkins et al. 1998). We consider four members of the CDM family: an open model (OCDM), a zero curvature low- Ω model (Λ CDM), and two models with an Einstein-de Sitter metric – the ‘standard CDM’ model (SCDM) and its modified version (τ CDM). Following Jenkins et al., the values, assigned to parameters $(h, \Omega, \Omega_\Lambda, \sigma_8)$ are: (0.7, 0.3, 0, 0.85) for OCDM, (0.7, 0.3, 0.7, 0.9) for Λ CDM, and (0.5, 1, 0, 0.6) for the SCDM and τ CDM which has extra large-scale power (added in an ad hoc manner, described by Jenkins et al. 1998). Here and below σ_8 is the rms dark matter density contrast in a ball of radius $8 h^{-1}$ Mpc.

Since CDM-like models are not scale-free, eq. (10) does not apply. In principle, we should therefore calculate $\alpha(x) \equiv \bar{\xi}^{(2)}(x, a)/[\bar{\xi}^{(1)}(x, a)]^2$ for each considered power spectrum and each separation x , using standard perturbative techniques (Lokas et al. 1996, Scoccimarro & Frieman 1996). However, as we will show below, these calculations can be significantly simplified by finding an effective slope, γ_{eff} , which provides ‘best fit’ α and v_{12} when substituted in eq. (10) and (11). The precise value of α is unimportant in the stable clustering regime as well as in the linear regime, when the term quadratic in $\bar{\xi}$ is sub-dominant. Hence, the precision in $\alpha(x)$ matters only at the boundary between the linear and nonlinear regimes, say at $\xi(x, t) = 1$. One of the possible definitions of the effective slope is therefore given by

$$\gamma_{\text{eff}} = -(d \ln \xi / d \ln x)|_{\xi=1}. \quad (13)$$

One can also choose γ_{eff} as follows. In Fig. 1, we plot logarithmic slopes of $\xi^{(1)}(x)$ and $\xi(x)$ (the latter is measured from the simulations). Both curves agree at large separations as they should, apart from small differences arising from noise in the measurement (we use only a finite number of bins and pairs to measure ξ), and from finite box-size and cosmic-variance effects (Jenkins et al. 1998). However, there is a well-defined scale (marked with an asterisk) at which the non-linear slope turns away from the linear theory prediction, marking the onset of the non-linear regime. We take γ_{eff} to be the logarithmic slope of $\xi^{(1)}$ at that scale. The resulting slopes are: $\gamma_{\text{eff}} = 1.67$ (SCDM), 1.46 (Λ CDM), 1.49 (OCDM), and 1.40 (τ CDM). The alternative definition (13) gives, respectively: $\gamma_{\text{eff}} = 1.67, 1.47, 1.45$, and 1.28. The advantage of the former definition is that it is more closely related to observations because it uses $\xi(x)$ only rather than ξ along with $\xi^{(1)}(x)$. The linear correlation function is easy to calculate under the controlled conditions of an N-body experiment but it can not be easily determined from observations.

In Figure 2 we test our Ansatz (11) against N-body measurements. For comparison, we also plot three other approximations for $v_{12}(r)$, considered earlier in the literature:

$$\text{A) } v_{12} = -\frac{2}{3}Hrf\bar{\xi}^{(1)} \quad , \quad \text{B) } v_{12} = -\frac{2}{3}Hrf\bar{\xi} \quad , \quad \text{C) } v_{12} = -\frac{2}{3}Hrf\bar{\xi} \quad . \quad (14)$$

(A) and (B) are two variants of linear theory predictions, and (C) is an improvement over linear theory, suggested by Peebles (LSS, §71). Figure 2 shows that the deviations from linear theory are small at large separations, as they should. The range of validity of the Peebles formula (C) is already considerably wider than that of linear theory. However, our new ansatz provides by far the best approximation. In fact, it covers the entire dynamical range!

The scale at which the linear approximation becomes acceptable depends on the amplitude of fluctuations; it increases with increasing σ_8 . For example, for a power-law correlation function, we have

$$\bar{\xi}(r) = \sigma_8^2 F(\gamma) r^{-\gamma} \quad , \quad \text{where} \quad (15)$$

$$F(\gamma) = (16 h^{-1} \text{Mpc})^\gamma (4 - \gamma)(6 - \gamma)/24 \quad , \quad (16)$$

and eq. (14 B) can be rewritten as

$$v_{12}(r) \approx -\frac{2}{3}\sigma_8^2 \Omega^{0.6} H F(\gamma) r^{1-\gamma} \approx -667 \sigma_8^2 \Omega^{0.6} \text{ km/s} \quad , \quad (17)$$

where the expression after the last “ $\approx$ ” sign assumes $\gamma = 1.75$ and $r = 10 h^{-1} \text{Mpc}$. The relative error in the latter expression, introduced by linear theory, can be calculated from eqs. (11) and (10). It depends on σ_8 only; for $\sigma_8 = 1$ and 0.6, linear theory overestimates $|v_{12}|$ by 24% and 10%, respectively.

3. Velocity bias

So far we considered the dynamics of pairwise motions of dark matter particles. However, for practical applications, it is necessary to understand the relation between $v_{12}(r)$ and the relative pairwise velocity of galaxies, $v_{12g}(r)$. We define the galaxy clustering bias as the square root of the ratio between the galaxy and the dark matter correlation functions: $b(r, t)^2 = \xi_g(r, t)/\xi(r, t)$.

Observations suggest that there is no velocity bias: splitting the Mark III catalogue (Willick et al. 1997) into subsamples of elliptical and spiral galaxies has no effect on v_{12g} (Juszkiewicz et al. 1998). Similar results follow from recent numerical simulations, which account for dissipative processes, important for galaxy formation. These models do show some *clustering bias* on small scales; however, there is no *velocity bias*, and $v_{12} = v_{12g}$ (Kauffmann et al. 1998).

As we show now, this contradicts the simplest toy model for bias, where $b \approx 1/\sigma_8$ is a constant. In this prescription, also known as linear bias, the galaxy density contrast at position $\vec{r}_A$ is simply given by $\delta_{gA} \approx b\delta_A$, where $\delta_A \equiv \rho_A/\langle\rho\rangle - 1$ is the mass density fluctuation, and $A = 1, 2, \dots$ enumerate galaxy positions. In the fluid approach, the mean pairwise velocity between two points at distance r is given by

$$\vec{v}_{12}(r) = \frac{\langle(\vec{v}_1 - \vec{v}_2)(1 + \delta_1)(1 + \delta_2)\rangle}{1 + \xi(r)}, \quad (18)$$

where $\vec{v}_A$ is the peculiar velocity at a point $\vec{r}_A$, $r = |\vec{r}_1 - \vec{r}_2|$ is the separation, and $\xi(r) = \langle\delta_1\delta_2\rangle$. For the galaxy pair density-weighted relative velocity, v_{12g} , the matter density field in the above expression, δ , has to be replaced by δ_g . In the limit of large separations ($\delta, \xi \rightarrow 0$), the linear biasing model, applied to eq. (18), gives $v_{12g}(r) = bv_{12}(r)$, and since $v_{12} \propto \sigma_8^2\Omega^{0.6}$ one obtains $v_{12g} \propto \sigma_8\Omega^{0.6}$ (Fisher et al. 1994). On small scales, where $1 + \delta \sim \delta$ and $1 + \xi \sim \xi$, the factors of b in the denominator and numerator cancel and $v_{12g}(r) = v_{12}(r)$. This unphysical behavior shows the limitations of the linear biasing model. Similar results can be obtained from considerations, based on the continuity equation and gravitational clustering. Gravity tends to remove bias and b has to evolve with time and separation (Fry 1996). The results of Kauffmann et al. (1998) fit neatly into this picture: their measurements of $b(r)$ at the ‘present time’ (redshift = 0) are significantly different from unity only on small scales where $\xi \gg 1$ and gravity is no longer the only force determining the dynamics; at large scales, however, gravity dominates and $b = 1$. Since the dependence on b in eq. (18) cancels out when $\xi \gg 1$, it is not surprising that the simulations give $v_{12} = v_{12g}$ on all scales. Summarizing, we conclude that a realistic model of biasing should lead to unbiased pairwise velocities; the linear bias model disagrees with observations as well as with simple theoretical considerations.

4. Conclusions

The main result of this paper is the expression (11) – an approximate analytic solution of the pair conservation equation. Our equations (10) - (13) reproduce the results of numerical simulations on all scales from the strongly non-linear to the linear regime.

Our results can be used to estimate Ω from redshift-distance surveys, like Mark III (Ferreira et al. 1998, Juszkiewicz et al. 1998). On very large scales, $v_{12}(r)$ is proportional to $\Omega^{0.6}\sigma_8^2$. On intermediate scales (the mildly non-linear regime) this degeneracy is removed and Ω and σ_8 can be measured separately because the $\alpha\bar{\xi}$ term in eq. (11) is Ω -independent. These properties make our estimator complementary to other estimators. Indeed, the POTENT method (Sigad et al. 1998) and the cluster abundances (Bahcall & Fan, 1998, Eke et al. 1998) are sensitive to $\beta \equiv \Omega^{0.6}\sigma_8$; the supernovae (Riess et al. 1998, Perlmutter et al. 1998) distances measure $q_o \equiv (\Omega/2) - \Omega_\Lambda$; finally, the position of acoustic peaks in the CMB power spectrum (Doroshkevich et al. 1978) is sensitive

to $\Omega + \Omega_\Lambda$. The advantage of our estimator over the CMB peaks method is model-independence; its advantage over POTENT is simplicity (Ferreira et al. 1998).

Our results can also be used to study the nature of biasing and to test the gravitational instability theory. Indeed, one can use the galaxy correlation function, $\xi_g(r)$, estimated from galaxy redshift or angular surveys to predict $v_{12}(r)$ which could then be compared to a $v_{12}(r)$ estimated from redshift-distance surveys (Gaztañaga & Juskiewicz 1998).

We thank Antonaldo Diaferio, Enrique Gaztañaga and Roman Scoccimarro for insightful comments. This work was supported by grants from the Polish Government (KBN grants No. 2.P03D.008.13 and 2.P03D.004.13), the Tomalla Foundation, by the Poland-US M. Skłodowska-Curie Fund and by the Swiss National Science Foundation. RJ thanks Simon White for his hospitality at MPA in Garching.

REFERENCES

- Bahcall, N.A. & Fan, X. 1998 ApJ, 504, 1
- Baugh, C.M., & Efstathiou, G. 1994, MNRAS, 270, 183
- Bouchet, F.R., Colombi, S., Hivon, E., & Juskiewicz, R. 1995, A&A, 296, 575
- Davis, M. & Peebles, P.J.E 1977, ApJS, 267, 465
- Doroshkevich, A.G., Zel’dovich, Ya.B. & Sunyaev, R.A. 1978, Sov.Astron., 22, 523.
- Ferreira, P.G., Juskiewicz, R., Feldman, H.A., Davis, M., & Jaffe, A.H. 1998, ApJ, in press
- Eke, V.R., Cole, S., Frenk, C.S. & Patrick, H.J., 1998 MNRAS, 298, 1145
- Fisher, K.B., et al. 1994, MNRAS, 267, 927
- Fry, J. 1984, ApJ, 279, 499
- Fry, J. 1996, ApJ, 461, L65
- Gaztañaga, E., Juskiewicz, R. 1998, in preparation
- Jain, B., Mo, H.J., & White, S.D.M. 1995, MNRAS, 276, L25
- Jenkins, A., et al. 1998, ApJ, 499, 20
- Juskiewicz, R. 1981, MNRAS, 197, 931
- Juskiewicz, R., Sonoda, D.H., & Barrow, J.D. 1984, MNRAS, 209, 139
- Juskiewicz, R., Bouchet, F.R., & Colombi, S. 1993, ApJ, 412, L9
- Juskiewicz, R., Davis, M., Feldman, H.A., Ferreira, P.G., Jaffe, A.H. 1998, in preparation

- Kauffmann, G., Colberg, J.M., Diaferio, A., & White, S.D.M. 1998, [astro-ph/9805283](#)
- Lokas, E., Juskiewicz, R., Bouchet, F., & Hivon, E. 1996, *ApJ*, 467, 1
- Mo, H.J., Jing, Y.P., & Börner, G. 1997, *MNRAS*, 286, 979
- Peebles, P.J.E. 1980, *The Large-Scale Structure of the Universe*, (Princeton: Princeton University Press) (LSS)
- Peebles, P.J.E. 1993, *Principles of Physical Cosmology*, (Princeton: Princeton University Press)
- Perlmutter, S. et al. 1998, *Nature*, 391, 51
- Riess, A. et al. 1998, *ApJ*, in press ([astro-ph/9805201](#))
- Scoccimarro, R., & Frieman, J. 1996, *ApJ*, 473, 620
- Sigad, Y. et al. 1998, *ApJ*, 495, 516
- Vishniac, E. 1983, *MNRAS*, 203, 345
- Willick, J.A., et al. 1997, *ApJS*, 109, 333




Fig. 1.— Logarithmic slope $d \ln \xi / d \ln r$ of the linear theory correlation function (dashed), and the measured non-linear correlation function (solid) for the four Virgo simulations which we have analyzed. The asterisks mark the effective slopes γ_{eff} used in equations (10) and (11), respectively.




Fig. 2.— The mean pairwise velocity v_{12} of the Virgo simulations (solid lines) compared with four closed-form approximations to solutions of the pair conservation equation: the two versions of the linear approximation, eq. 14 (A), and (B), are plotted as thin and thick dashed curves respectively; the Peebles approximation, eq. 14 (C), is shown as dot-dashed curve; and eq. (11) – the Ansatz proposed in this paper – is drawn as dot-dot-dot-dashed line.